

Classical Mechanics Phy 235, Lecture 11.

Frank L. H. Wolfs
Department of Physics and Astronomy
University of Rochester

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 1

Hamilton's Principle

"Of all the possible paths along which a dynamical system may move from one point to another within a specified time interval (consistent with any constraints), the actual path followed is that which minimizes the time integral of the difference between the kinetic and potential energies."

$$\delta \int_{t_1}^{t_2} (T - U) dt = 0$$

The quantity $T - U$ is called the **Lagrangian L** .

The Lagrange equation(s) of motion are:

$$\frac{\partial L}{\partial x_i} - \frac{d}{dt} \frac{\partial L}{\partial \dot{x}_i} = 0$$

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 2

Problem 7.12

A particle of mass m rests on a smooth plane.

The plane is raised to an inclination angle θ at a constant rate a ($\theta = 0^\circ$ at $t = 0$ s), causing the particle to move down the plane.

Determine the motion of the particle.

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 3

Generalized coordinates.

- So far we have expressed the Lagrangian in terms of (generalized) position and (generalized) velocities:

$$L = T - U = \frac{1}{2} m \sum_{i=1}^3 \dot{x}_i^2 - U(x_i)$$

- An alternative is to express the Lagrangian in terms of (generalized) position and (generalized) momenta. For example:

$$p_r = \frac{\partial L}{\partial \dot{r}} = \frac{\partial T}{\partial \dot{r}} = m\dot{r}$$

$$p_\theta = \frac{\partial L}{\partial \dot{\theta}} = \frac{\partial T}{\partial \dot{\theta}} = m r^2 \dot{\theta}$$

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 4

Conservation Laws – Part I.

- Conservation of energy:
 - Lagrangian does not depend on time explicitly.
 - If L does not depend explicitly on time, it can be shown that

$$L - \sum_j \left(\dot{q}_j \frac{\partial L}{\partial \dot{q}_j} \right) = \text{constant} = -H$$

- The constant H is called the Hamiltonian of the system:

$$H = \sum_j \left(\dot{q}_j \frac{\partial L}{\partial \dot{q}_j} \right) - L$$

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 5

Problem 7.22.

A particle of mass m moves in one dimension under the influence of a force F :

$$F(x,t) = \frac{k}{x^2} e^{-t/\tau}$$

where k and τ are positive constants. Compute the Lagrangian and Hamiltonian functions. Compare the Hamiltonian and the total energy and discuss the conservation of energy for the system.

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 6



2 Minute 2 Second Intermission.

- Since paying attention for 1 hour and 15 minutes is hard when the topic is physics, let's take a 2 minute 2 second intermission.
- You can:
 - Stretch out.
 - Talk to your neighbors.
 - Ask me a quick question.
 - Enjoy the fantastic music.



Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 7

Conservation Laws – Part II.

- Conservation of linear momentum:
 - Lagrangian should not be effected by a translation of space.
- Conservation of angular momentum:
 - Lagrangian should not be effected by a rotation of space.

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 8

Canonical Equations of Motion

Lagrange equations of motion in terms of generalized momentum:

$$\frac{d}{dt} \frac{\partial L}{\partial \dot{q}_i} = \dot{p}_i = \frac{\partial L}{\partial q_i}$$

The Hamiltonian H can be written in terms of the generalized momenta as

$$H = \sum_j \dot{q}_j \frac{\partial L}{\partial \dot{q}_j} - L = \sum_j \dot{q}_j p_j - L$$

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 9

Hamilton's Equations of Motion

- For each coordinate: **two** equations of motion.
 - For each coordinate there is only one Lagrange equation of motion.
- Equations of motion are **first order differential equations**.
 - The Lagrange equations of motion are second order differential equations.

$$\frac{\partial H}{\partial p_j} - \dot{q}_j = 0$$

$$\frac{\partial H}{\partial q_j} + \dot{p}_j = 0$$

$$\frac{\partial H}{\partial t} + \frac{\partial L}{\partial t} = 0$$

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 10

Problem 7.38

The potential for an anharmonic oscillator is $U = kx^2/2 + bx^4/4$ where k and b are constants.

Find Hamilton's equations of motion.

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 11

Problem 7.28

The force F that is provided fixed the potential U :

$$U = -\frac{k}{r}$$

Use plane polar coordinates and find Hamilton's equations of motion.

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 12

Problem 7.24

Consider a simple plane pendulum consisting of a mass m attached to a string of length l . After the pendulum is set into motion, the length of the string is shortened at a constant rate:

$$\frac{dl}{dt} = -\alpha = \text{constant}$$

The suspension point remains fixed. Compute the Lagrangian and Hamiltonian functions. Compare the Hamiltonian and the total energy, and discuss the conservation of energy for the system.

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 13

End of Chapter 7

- We now have finished the material that will be covered on midterm exam# 2: chapters 5 – 7.
- Midterm exam # 2 will take place on Tuesday October 23 between 8 am and 9.30 am in B&L 109.

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 14

ENOUGH FOR TODAY?

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 15
