

Classical Mechanics
Phy 235, Lecture 16.

Frank L. H. Wolfs
Department of Physics and Astronomy
University of Rochester

Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester, Slide 1

A few comments

- On the TA/TAI survey, a few students stated that certain characters do not display correctly on the PHY 235 website. This is a limitation of HTML, but all documents (notes and homework assignments) are also available from the download page, where symbols are displayed properly. See: <http://teacher.pas.rochester.edu/PHY235/DownloadFolder/DownloadIndex.htm>
- On the TA/TAI survey, a student wrote “his no curve policy seems like a ploy to fail us all”. I respectfully disagree. Looking at last years record, no one failed PHY 235 and the average grade was a B+. Grading on a curve would have lowered everyones grade since a B average is commonly used for curving.

Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester, Slide 2

Motion in Non-Inertial Reference Frames.

- The Earth is not an inertial reference frame:
 - The Earth is rotating around its axis.
 - The Earth is moving around the sun.
 - The sun is moving around the center of the Milky-Way galaxy.
- What is the effect of this non-inertial motion?
- All will be revealed in Chapter 10.

Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester, Slide 3

Problem 10.1.


Calculate the centrifugal acceleration due to the Earth's rotation on a particle on the surface of the Earth at the equator. Compare this result with the gravitational acceleration.

Compute also the centrifugal acceleration due to the motion of the Earth about the sun, and the motion of the sun around the center of the Milky Way.

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 4

Rotating Coordinate system.



Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 5

The angular acceleration is the same in both reference frames.

- Relation between position vectors:

$$\left(\frac{d\vec{r}}{dt}\right)_{fixed} = \left(\frac{d\vec{r}}{dt}\right)_{rotating} + \vec{\omega} \times \vec{r}$$

- Relation between angular velocity vectors:

$$\left(\frac{d\vec{\omega}}{dt}\right)_{fixed} = \left(\frac{d\vec{\omega}}{dt}\right)_{rotating} + \vec{\omega} \times \vec{\omega} = \left(\frac{d\vec{\omega}}{dt}\right)_{rotating}$$

- Conclusion: the angular acceleration is the same in both reference frames.

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 6

Velocity in fixed (inertial) frame.

Velocity of the origin of the rotating frame.


$$v_f = \left(\frac{d\vec{r}'}{dt} \right)_{\text{fixed}} = \left(\frac{d\vec{R}}{dt} \right)_{\text{fixed}} + \left(\frac{d\vec{r}}{dt} \right)_{\text{rotating}} + \vec{\omega} \times \vec{r} = V + v_r + \vec{\omega} \times \vec{r}$$

Velocity in fixed frame. Velocity in rotating frame.

Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester, Slide 7

3 Minute 10 Second Intermission.

- Since paying attention for 1 hour and 15 minutes is hard when the topic is physics, let's take a 3 minute 10 second intermission.
- You can:
 - Stretch out.
 - Talk to your neighbors.
 - Ask me a quick question.
 - Enjoy the fantastic music.



Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester, Slide 8

Newton's laws in rotating reference frames.

- Only in the fixed reference frame can we use Newton's second law:

$$\vec{a}_f = \left(\frac{d\vec{v}_f}{dt} \right)_{\text{fixed}} = \frac{\vec{F}}{m}$$
- The acceleration in the fixed reference frame can also be expressed as:

$$\vec{a}_f = \left(\frac{d\vec{V}}{dt} \right)_{\text{fixed}} + \left(\frac{d\vec{v}_r}{dt} \right)_{\text{rotating}} + 2\vec{\omega} \times \vec{v}_r + \dot{\vec{\omega}} \times \vec{r} + \vec{\omega} \times \{ \vec{\omega} \times \vec{r} \}$$

Acceleration observed in rotating frame.

Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester, Slide 9

Observed acceleration in rotating reference frame.

- An observer in the rotating reference frame will observe an acceleration:

$$\vec{a}_r = \left(\frac{d\vec{v}}{dt} \right)_{\text{rotating}}$$

- This acceleration can be expressed in terms of the acceleration in the inertial frame:

$$\vec{a}_r = \vec{a}_f - \left(\frac{d\vec{v}}{dt} \right)_{\text{fixed}} - 2\vec{\omega} \times \vec{v}_r - \dot{\vec{\omega}} \times \vec{r} - \vec{\omega} \times \{ \vec{\omega} \times \vec{r} \}$$

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 10

The effective force.

Using the **effective force**

$$\vec{F}_{\text{eff}} = m\vec{a}_f - m\dot{\vec{\omega}} \times \vec{r} - 2m\vec{\omega} \times \vec{v}_r - m\vec{\omega} \times \{ \vec{\omega} \times \vec{r} \}$$

Coriolis force.

Centripetal force.



Not Dutch!
So forget it.

an observer in the rotating frame will be able to determine the acceleration in the rotating frame by dividing the effective force by the mass of the object.

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 11

Problem 10-6.

A bucket of water is set spinning about its symmetry axis. Determine the shape of the water in the bucket.

Frank L. H. Wolfs

Department of Physics and Astronomy, University of Rochester, Slide 12

ENOUGH FOR TODAY?

Frank L. H. Wolfs Department of Physics and Astronomy, University of Rochester, Slide 13
